

Continuation of the MOLLUSCA of the Chevert Expedition, by  
J. BRAZIER, C.M.Z.S., Corr. Memb. Roy. Soc., Tas.

#### SUB FAMILY FUSINÆ.

##### 1.—FUSUS PROBOSCIDIFERUS.

*Fusus proboscidiferus*, Lam. Anim. Sans, Vert., tome 7, p. 126.

“ “ Lam. (Deshayes) edit., vol. 9, p. 447.

“ “ Reeve, Conch. Icon., pl. 4, sp. 15.

“ “ Chenu. Manuel de Conch., p. 139, fig. 596.

*Hab.* Darnley Island, Torres Straits. Found on the outer reefs, specimen from 18 to 24 inches long.

#### FAMILY PLEUROTOMIDÆ.

##### SUB FAMILY PLEUROTOMINÆ.

##### 2.—PLEUROTOMA SPECTABILIS.

*Pleurotoma spectabilis*, Reeve, Conch. Icon., pl. 1, sp. 6.

*Hab.* Darnley Island, Torres Straits, 20 fathoms, sand.

##### 3.—PLEUROTOMA TIGRINA.

*Pleurotoma tigrina*, Lam. Anim. Sans, Vert., tome 7, p. 95.

“ *marmorata*, Lam. Encyclop., pl. 439, f. 6.

“ *tigrina*, Reeve, Conch. Icon., pl. 1, sp. 3.

*Hab.* Princess Charlotte Bay, North East Australia, 14 fathoms; Darnley Island, Torres Straits, 30 fathoms, sand bottom.

##### 4.—PLEUROTOMA CRISPA.

*Pleurotoma crispa*, Lam. Anim. Sans, Vert., tome 7, p. 95.

“ “ Reeve, Conch. Icon., pl. 2, sp. 11 a, b.

*Hab.* Darnley Island, Torres Straits, 20 fathoms, sandy mud.

##### 5.—PLEUROTOMA MARMORATA.

*Pleurotoma marmorata*, Lam. Anim. Sans, Vert., tome 7, p. 95.

“ “ Var. *maculata*, Reeve, Conch. Icon., sp. 21a.

*Hab.* Darnley Island, Torres Straits, 25 fathoms.

#### FAMILY TEREBRIDA.

##### 6.—ACUS MUSCARIA.

*Terebra muscaria*, Lam. Anim. Sans, Vert., tome 7, p. 285.

“ *subulata*, Lam. Encyclop., pl. 402, f. 2a, b.

“ *muscaria*, Sowerby, Thes. Conch., vol. 1, p. 154, pl. 42, fig. 41.

*Hab.* Darnley Island, Torres Straits. Found on the sands inside the reef.

7.—**ACUS SUBULATA.**

*Buccinum subulatum*, Linn. Syst. Nat., edit. 12, p. 1205.

*Terebra subulata*, Lam. Anim. Sans, Vert., tome 7, p. 242.

“ “ Sowerby, Thes. Conch., vol. 1, p. 156, pl. 41,  
fig. 16.

*Buccinum subulatum*, Hanley, Ipsa, Linnæi, Conchylia, p. 259.

*Hab.* Darnley Island, Torres Straits. Found on the reefs inside the reefs.

**FAMILY CERITHIIDÆ.**

8.—**CERITHIUM (COLINA) COARCTATUM.**

*Cerithium coarctatum*, Sowerby, Thes. Conch., vol. 3, Thes., pl. 290, fig. 321, 322.

*Hab.* Bet Island, Torres Straits, 11 fathoms, coral and sand, very rare, only one specimen found.

**FAMILY CAPULIDÆ.**

9.—**HIPPONYX BARBATA.**

*Hipponyx barbata*, Sowerby, Proc. Zool. Soc., 1835, p. 5; Thes. Conch., vol. 1, p. 369, pl. 73, fig. 26, 27.

*Hab.* Cape York, North Australia, 5 fathoms, one dead specimen brought up in the dredge. Anse Vata, near Noumea, New Caledonia. (Brazier.)

Notes and remarks on MOLLUSCA recently found in Port Jackson and New Caledonia, by J. BRAZIER, C.M.Z.S., Corr. Mem. Roy. Soc., Tas.

1.—**HALIOTIS PARVA.**

*Haliotis parva*, Linn. Syst. Nat. ed. 12, p. 1256.

“ “ *canaliculata*, Lam. Anim. Sans, Vert., tome 6, second part, p. 217.

*Haliotis parva*, Reeve, Conch. Icon., pl. 15, fig. 53b.